

Supersymmetric strictly isospectral FRW models for zero factor ordering

H.C. Rosu and J. Socorro

Instituto de Física de la Universidad de Guanajuato, Apdo Postal E-143, León, Gto, México

We apply the strictly isospectral technique of standard supersymmetric quantum mechanics to the $Q = 0$ factor ordered Wheeler-DeWitt equation for the closed Friedmann-Robertson-Walker (FRW) minisuperspace model. The resulting strictly isospectral one-parameter families of both FRW cosmological potentials and “wavefunctions of the universe” are exhibited with relevant plots.

PACS numbers: 98.80.Hw, 11.30.Pb, 04.60.Kz

gr-qc/9606030

Nuovo Cimento B 113, 683-689 (1998)

Bianchi cosmologies [1] are cosmological models which are currently used to illustrate possible ways of finding exact solutions to the canonical constraints in quantum cosmology. In the last decade the number of exact solutions has been continuously increased by applying various formalisms [2]. The simplest of all cosmologies is the FRW model because of its full one-dimensional character. The goal of the present work is to apply the strictly isospectral scheme based on the general Riccati solution [3,4], which is quite well known in nonrelativistic one-dimensional supersymmetric quantum mechanics, to the $Q = 0$ factor-ordered Wheeler-DeWitt (WDW) equation corresponding to the closed FRW cosmological model. Thus, we will be able to display a one-parameter class of strictly isospectral cosmological FRW solutions and to present the wavefunctions of the universe for that case.

FRW models belongs to the diagonal Bianchi type IX models in the class A models [1]. Therefore we recall briefly the canonical formulation in the Arnowitt-Deser-Misner (ADM) formalism of the latter ones. The metric has the form

$$ds^2 = -dt^2 + e^{2\alpha(t)} (e^{2\beta(t)})_{ij} \omega^i \omega^j, \quad (1)$$

where $\alpha(t)$ is a scalar function and $\beta_{ij}(t)$ a 3x3 diagonal matrix, $\beta_{ij} = \text{diag}(x + \sqrt{3}y, x - \sqrt{3}y, -2x)$, ω^i are one-forms which characterize each cosmological Bianchi type model, obeying $d\omega^i = \frac{1}{2}C_{jk}^i \omega^j \wedge \omega^k$, C_{jk}^i being the structure constants of the corresponding invariance group.

The ADM action has the form

$$I = \int (P_x dx + P_y dy + P_\alpha d\alpha - N \mathcal{H}_\perp) dt, \quad (2)$$

where the P 's are canonically conjugated momenta, N is the lapse function and

$$\mathcal{H}_\perp = e^{-3\alpha} (-P_\alpha^2 + P_x^2 + P_y^2 + e^{4\alpha} V(x, y)) . \quad (3)$$

$e^{4\alpha} V(x, y) = U(q^\mu)$ is the potential term of the cosmological model under consideration. The WDW equation for these models is achieved by replacing P_{q^μ} by $-i\partial_{q^\mu}$

in Eq. (3), with $q^\mu = (\alpha, x, y)$. The factor $e^{-3\alpha}$ may be factor-ordered with P_{q^μ} in many ways. Hartle and Hawking [5] have suggested an almost general factor ordering, which in this case would order $e^{-3\alpha} P_{q^\mu}^2$ as

$$-e^{-(3-Q)\alpha} \partial_\alpha e^{-Q\alpha} \partial_\alpha = -e^{-3\alpha} \partial_\alpha^2 + Q e^{-3\alpha} \partial_\alpha, \quad (4)$$

where Q is any real constant. In the following, we shall work in the $Q = 0$ factor ordering in which the WDW equation reads

$$\square \Psi - U(q^\mu) \Psi = 0, \quad (5)$$

and by means of the Ansatz $\Psi(q^\mu) = A e^{\pm\Phi}$ one gets

$$\pm A \square \Phi + A[(\nabla\Phi)^2 - U] = 0, \quad (6)$$

where $\square = G^{\mu\nu} \frac{\partial^2}{\partial q^\mu \partial q^\nu}$, $(\nabla)^2 = -(\frac{\partial}{\partial \alpha})^2 + (\frac{\partial}{\partial x})^2 + (\frac{\partial}{\partial y})^2$, and $G^{\mu\nu} = \text{diag}(-1, 1, 1)$.

For the FRW model there is only the α coordinate and the corresponding potential is $U = -e^{4\alpha}$. Then the WDW equation can be written as a one-dimensional ordinary differential equation

$$\left(\frac{d\Phi}{d\alpha}\right)^2 \pm \left(\frac{d^2\Phi}{d\alpha^2}\right) = e^{4\alpha} \quad (7)$$

and allows one to think of the Witten supersymmetric procedure (for a recent review see [6]) since the change of function $\Phi = \pm \ln y$ produces a zero-energy one-dimensional Schrödinger equation, whereas the change $\frac{\partial\Phi}{\partial\alpha} = \pm\mathcal{W}$ leads to a Riccati equation for $\mathcal{W}$, which can be identified with the superpotential in the Witten procedure. The one-dimensional Schrödinger equation reads

$$-y'' + e^{4\alpha} y = 0, \quad (8)$$

with the second derivative taken with respect to the α coordinate, and possessing as general solution a superposition of modified Bessel functions $Z_0(i\frac{1}{2}e^{2\alpha}) = C_1 I_0(\frac{1}{2}e^{2\alpha}) + C_2 K_0(\frac{1}{2}e^{2\alpha})$. Let us consider separately the two particular cases as follows:

(i) $Z_0 = K_0$; This case corresponds asymptotically to an Ansatz of the type $\Psi = Ae^{-\Phi_1}$ for the “wavefunction of the universe”, with $\Phi_1 = \frac{1}{2}e^{2\alpha} + \alpha$. Indeed, asymptotically ($\alpha \gg 1$) [7]

$$K_0(\frac{1}{2}e^{2\alpha}) \approx e^{-\frac{1}{2}e^{2\alpha} - \alpha}. \quad (9)$$

(ii) $Z_0 = I_0$; Again asymptotically, it corresponds to an Ansatz of the type $\Psi = Ae^{+\Phi_2}$ for the ‘wavefunction of the universe’, with $\Phi_2 = \frac{1}{2}e^{2\alpha} - \alpha$ since for $\alpha \gg 1$ [7]

$$I_0(\frac{1}{2}e^{2\alpha}) \approx e^{\frac{1}{2}e^{2\alpha} - \alpha}. \quad (10)$$

We shall now apply the technique of strictly isospectral potentials to the FRW model. This technique is known since about a decade in one-dimensional supersymmetric quantum mechanics and usually requires nodeless, normalizable states of Schrödinger equation. However, Pappademos, Sukhatme, and Pagnamenta [8] showed that the strictly isospectral construction can be also performed on nonnormalizable states. Thus, using the nonnormalizable modified Bessel function Z_0 the strictly isospectral construction says that the isospectral potentials for the FRW cosmological model are given by

$$\begin{aligned} S_{iso}(\alpha; \lambda) &= S^-(\alpha) - 2[\ln(\mathcal{J}_0 + \lambda)]'' \\ &= S^-(\alpha) - \frac{4Z_0(\alpha)Z_0'(\alpha)}{\mathcal{J}_0 + \lambda} + \frac{2Z_0^4(\alpha)}{(\mathcal{J}_0 + \lambda)^2}, \end{aligned} \quad (11)$$

where $S^-(\alpha) = e^{4\alpha}$ is the Schrödinger-FRW potential, λ is a real parameter and

$$\mathcal{J}_0(\alpha) \equiv \int_{-\infty}^{\alpha} Z_0^2(i\frac{1}{2}e^{2y})dy. \quad (12)$$

The common argument for the supersymmetric strictly isospectral method is as follows. One can factorize the FRW one-dimensional Schrödinger equation with the operators $A = \frac{d}{d\alpha} + \mathcal{W}(\alpha)$ and $A^\dagger = -\frac{d}{d\alpha} + \mathcal{W}(\alpha)$, where the superpotential function is given by $\mathcal{W} = -Z_0'/Z_0$. Then, in the Witten scheme, the potential and superpotential enter an initial ‘bosonic’ Riccati equation $S^- = \mathcal{W}^2 - \mathcal{W}'$. At the same time, one can build a ‘fermionic’ Riccati equation $S^+ = \mathcal{W}^2 + \mathcal{W}'$, corresponding to a ‘fermionic’ Schrödinger equation for which the operators A and $A^\dagger$ are applied in reversed order. Thus, the ‘fermionic’ Schrödinger-FRW potential is found to be $S^+ = e^{4\alpha} - 2(\frac{Z_0'}{Z_0})'$. This potential does not have the Z_0 solution as a ‘wavefunction of the universe’. However, it is possible to reintroduce the Z_0 solution into the spectrum, by using the general superpotential solution of the ‘fermionic’ Riccati equation. The general Riccati solution reads

$$\mathcal{W}_{gen} = \mathcal{W}(\alpha) + \frac{d}{d\alpha} \ln[\mathcal{J}_0(\alpha) + \lambda]. \quad (13)$$

The way to obtain Eq. (13) is well-known [3,4] and will not be repeated here. From it one can easily get Eq. (11). The one-parameter family of ‘wavefunctions of the universe’ differs from the initial one, being the λ -depending quotient

$$Z_{gen}(\alpha; \lambda) = \sqrt{\lambda(\lambda + 1)} \frac{Z_0(i\frac{1}{2}e^{2\alpha})}{\mathcal{J}_0 + \lambda}, \quad (14)$$

though in the limit $\lambda \rightarrow \pm\infty$ one recovers the initial solution. In Eq. (14) we have included the factor $\sqrt{\lambda(\lambda + 1)}$, however not for normalization reasons as in supersymmetric quantum mechanics [9] (the wavefunctions are still nonnormalizable) but in order to discuss the so-called Pursey limit [10] $\lambda = 0$, and the Abraham-Moses limit [11] $\lambda = -1$ (see figure captions). In order to get a better feeling of what is going on we have plotted the strictly isospectral FRW cosmological potentials and wavefunctions for $Z_0 = I_0$ and $Z_0 = I_0 - K_0$ and two values of the λ -parameter, see Figures 1, 2 and 3, confirming in this cosmological context some known facts of the supersymmetric strictly isospectral construction in nonrelativistic quantum mechanics [6]. What we actually did was to calculate the integral and all the other quantities in the interval $(-c, \alpha)$, where c is a sufficiently big positive number, while α has been chosen a not too big positive number. In other words, we have a particle (the FRW universe) in a special type of box, that we call a Bessel strictly isospectral box. For I_0 , the integral $\mathcal{J}_0$ is a finite quantity even in the infinite limit, and the computer calculations show small well structures in the rising branch of the isospectral potentials and corresponding peaks in the wavefunctions. The Bessel box is single-walled at the α limit. For $I_0 - K_0$, the integral is divergent in the infinite limit. In this case the Bessel strictly isospectral box is like having walls of different heights at the c limit and α limit. This peculiar box is a quite interesting object being the physical embodiment of the strictly isospectral family of FRW potentials as a family of Bessel solitonic potentials [12] possessing translationally moving structures. It can also be considered as a kind of cavity with slowly moving walls and wells if λ is varied even if the interval $(-c, \alpha)$ is kept fixed.

In conclusion, we showed that the Hamiltonian constraint of canonical quantum gravity for FRW minisuperspace models may contain a one-parameter family of FRW cosmological potentials and “wavefunctions of the universe”. They are strictly isospectral (i.e. at zero energy) WDW solutions of the FRW case including the common solution in the limit $\lambda \rightarrow \pm\infty$. Unfortunately, the strictly isospectral method can be applied only to either pure one-dimensional cases as the FRW model here (see also [13]) or to separable cases [14].

Acknowledgments

This work was supported in part by the CONACyT Projects 4868-E9406 and 3898P-E9608.

-
- [1] M. Ryan, *Hamiltonian Cosmology* (Springer, Berlin, 1972).
 - [2] O. Obregón and J. Socorro, Int. J. Theor. Phys. **35**, 1381 (1996) and references therein.
 - [3] B. Mielnik, J. Math. Phys. **25**, 3387 (1984).
 - [4] M.M. Nieto, Phys. Lett. B **145**, 208 (1984).
 - [5] J. Hartle and S.W. Hawking, Phys. Rev. D **28**, 2960 (1983).
 - [6] F. Cooper, A. Khare, and U. Sukhatme, Phys. Rep. **251**, 267 (1995).
 - [7] F. Bowman, *Introduction to Bessel Functions* (Dover Publ. Inc., 1958), formulas 5.45 and 5.44, respectively, where only the first leading term has been retained.
 - [8] J. Pappademos, U. Sukhatme, and A. Pagnamenta, Phys. Rev. A **48**, 3525 (1993).
 - [9] A. Khare, private communication.
 - [10] D. Pursey, Phys. Rev. D **33**, 1098, 1103, 2267 (1986); M. Luban and D. Pursey, Phys. Rev. D **33**, 431 (1986).
 - [11] P. Abraham and H. Moses, Phys. Rev. A **22**, 1333 (1980).
 - [12] W.-Y. Keung, U.P. Sukhatme, Q. Wang, and T.D. Imbo, J. Phys. A **22**, L987 (1989).
 - [13] H.C. Rosu and J. Socorro, Phys. Lett. A **223**, 28 (1996).
 - [14] H.C. Rosu and J. Socorro, Nuovo Cimento B **113**, 677 (1998) (gr-qc/9610018).




Fig. 1a

The original potential and two members of the one-parameter strictly isospectral family of FRW potentials in the case $Z_0 = I_0$ for $\lambda = 25$ (small well) and 100 (bigger well)




Fig. 1b
The corresponding wavefunctions.




Fig. 2a
The original potential and two members of the one-parameter strictly isospectral family of FRW potentials for $I_0 - K_0$, and the same values of the λ -parameter. On the arbitrary vertical scale the two peaks close at a value of about 2,000.




Fig. 2b

All three potentials on the same plot, showing at the left the peak corresponding to $\lambda = 100$ and slightly at the right that for $\lambda = 25$. We have found that the peaks are moving to the left if λ is positive and increasing (i.e., beyond Pursey limit), while they are moving toward the origin if λ is less than the Abraham-Moses limit and decreasing.




Fig. 2c

A more detailed plot of the region close to the origin for the three potentials showing the real trend of the plateau between the small well structures at the right and the peaks at the left.




Fig. 3

The strictly isospectral ‘wavefunction of the universe’, Z_{gen} , for $Z_0 = I_0 - K_0$ and the λ parameters as in Figs. 1 and 2 (the small peak corresponds to $\lambda = 25$, whereas the big one is for $\lambda = 100$). We notice also that the positive tails (at the right of the peaks) are practically zero, while the negative tails have a small positive slope. That implies that the wavefunctions are not normalizable. The segment very close to the vertical axis is a portion of the original ‘wavefunction’, close to the value $I_0(1/2) - K_0(1/2)$.

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/gr-qc/9606030v4>

This figure "fig1-2.png" is available in "png" format from:

<http://arXiv.org/ps/gr-qc/9606030v4>